

Table 1: Estimated impact of suppression strategies. The impact on infections and deaths over 250 days for two different suppression strategies triggered according to different mortality thresholds for mortality incidence (0.2 and 1.6 deaths per 100,000 population per week).

	Unmitigated Scenario		Suppression at 0.2 deaths per 100,000 population per week		Suppression at 1.6 deaths per 100,000 population per week	
	Infections	Deaths	Infections	Deaths	Infections	Deaths
East Asia & Pacific	2,117,131,000	15,303,000	92,544,000	442,000	632,619,000	3,315,000
Europe & Central Asia	801,770,000	7,276,000	61,578,000	279,000	257,706,000	1,397,000
Latin America & Caribbean	566,993,000	3,194,000	45,346,000	158,000	186,595,000	729,000
Middle East & North Africa	419,138,000	1,700,000	30,459,000	113,000	152,262,000	594,000
North America	326,079,000	2,981,000	17,730,000	92,000	90,529,000	520,000
South Asia	1,737,766,000	7,687,000	111,703,000	475,000	629,164,000	2,693,000
Sub-Saharan Africa	1,044,858,000	2,483,000	110,164,000	298,000	454,968,000	1,204,000
Total	7,013,734,000	40,624,000	469,523,000	1,858,000	2,403,843,000	10,452,000